








EMS System for Metropolitan Oklahoma City and Tulsa 2019 Medical Control Board Treatment Protocols



Approved 9/12/18, Effective 1/15/19, replaces all prior versions

1D - TRAUMA AND HYPOVOLEMIC SHOCK SUPPORTIVE CARE ADULT & PEDIATRIC

- TREATMENT PRIORITIES**
- Assessment:
 - SCENE SAFETY
 - PROTECTIVE EQUIPMENT
 - Primary Survey
 - "Trauma Alert" to receiving ED if indicated
 - Secondary Survey (when appropriate)
 - Primary Survey Care:
 - Control arterial bleeding
 - Open airway
 - Seal "sucking" chest wound(s)
 - Needle thoracostomy for closed chest tension pneumothorax
 - Minimize scene time in critical case.
 - Enroute Care:
 - Reassess all primary care
 - Support oxygenation/ventilation
 - Vascular access
 - Secondary Survey (if able)
 - Keep patient warm/avoid hypothermia
 - Hospital per destination protocol..



Clinical Operational Note (All Field Provider Levels): The practice of EMS medicine is built upon the foundation of "taking medical care to the patient". To achieve this objective, appropriate equipment (airway equipment kit, med/trauma equipment kit, suction device, patient packaging equipment) should be brought to the patient's side per Protocol 14J – Scene Coordination to minimize critical treatment delays