



EMS System for Metropolitan Oklahoma City and Tulsa 2019 Medical Control Board Treatment Protocols



Approved 9/12/18, Effective 1/15/19, replaces all prior versions

13D – COMPLICATIONS OF PREGNANCY ADULT

- TREATMENT PRIORITIES**
1. Vital Signs
 2. Dextrose for hypoglycemia
 3. Magnesium for eclampsia

EMD

ADVISE TO AVOID PHYSICAL EXERTION
OR ENVIRONMENTAL STRESS (TEMP EXTREMES).

EMERGENCY MEDICAL DISPATCHER
EMERGENCY MEDICAL RESPONDER
EMT
EMT-INTERMEDIATE 85
ADVANCED EMT
PARAMEDIC

EMR	EMT
GENERAL SUPPORTIVE CARE OBTAIN VITAL SIGNS O ₂ VIA NC or NRB AS INDICATED APPLY CARDIAC MONITOR/OBTAIN 12-LEAD ECG (if equipped) TRANSMIT 12-LEAD ECG TO RECEIVING EMERGENCY DEPARTMENT DETERMINE BLOOD GLUCOSE HYPOGLYCEMIA CARE: IF GLUCOSE <50 mg/dL, 1 tube ORAL GLUCOSE (15 grams) PO	

EMT-185	AEMT
IV ACCESS HYPOGLYCEMIA CARE: IF GLUCOSE <50 mg/dL, D50 1 mL/kg IVP UP TO 50 mL GLUCAGON 1 mg IM IF NO VASCULAR ACCESS OBTAINED	

PARAMEDIC

ECLAMPSIA: MAGNESIUM SULFATE 1 gram IVP/IOP
MAY REPEAT EVERY 2-3 MINUTES UNTIL SEIZURE ABATES
MAXIMUM CUMULATIVE DOSE IS 4 grams