



EMS System for Metropolitan Oklahoma City and Tulsa 2021 Medical Control Board Treatment Protocols



Approved 9/9/20, Effective 1/15/21, replaces all prior versions

16E – ASPIRIN

EMERGENCY MEDICAL DISPATCHER
EMERGENCY MEDICAL RESPONDER
EMT
EMT-INTERMEDIATE 85
ADVANCED EMT
PARAMEDIC

Self-Administration Phone Directive - 5A 5C

Assist Pt with Self Administration - 5A 5C

Class: Anti-Platelet

Actions/Pharmacodynamics: Inhibits platelet aggregation (and thereby, further clot formation). This action results in an overall increase in survival from acute myocardial infarction.

Indications: Chest Pain - Uncertain Etiology (5A)
Acute Coronary Syndrome (5C)

Contraindications: Active gastrointestinal bleeding
History of aspirin allergy including angioedema and/or anaphylaxis
History of asthma with aspirin-induced exacerbation

Pharmacokinetics: Absorption in stomach and small intestine, with onset of action within 30 minutes and duration of action for several hours.

Side Effects: Typically none from single EMS dosing. Rare instances of nausea or allergic reaction could be encountered. Treat allergic reaction per Protocol 8D - Acute Allergic Reactions.

Dosage: Chest Pain - Uncertain Etiology - Adult (5A)
Acute Coronary Syndrome - Adult (5C)
324 OR 325 mg chewed by patient (hold if taken 324+mg within 6 hours)

How Supplied: 81 mg tablets
325 mg tablets
(Always check concentration and dose per container at time of patient medication administration)

Special Comment: Aspirin is indicated even if the patient is taking warfarin sodium (Coumadin®), clopidogrel (Plavix®), or other anticoagulant or antiplatelet agents on a daily basis.