



EMS System for Metropolitan Oklahoma City and Tulsa 2025 Medical Control Board Treatment Protocols

Approved 9/04/24, Effective 1/15/25, replaces all prior versions

17F - MEDICATION-ASSISTED INTUBATION PEDIATRIC

TREATMENT PRIORITIES

1. Oxygenation/Ventilation support

EMERGENCY MEDICAL
DISPATCHER

EMERGENCY MEDICAL
RESPONDER

EMT

EMT-INTERMEDIATE 85

ADVANCED EMT

PARAMEDIC

PARAMEDIC

MEDICATION-ASSISTED INTUBATION IF INDICATED
FOLLOW PROTOCOL 2G – ORAL INTUBATION FOR TECHNIQUE & CONFIRMATION OF INTUBATION

FOR FACILITATING ORAL INTUBATION:

PEDIATRIC: ETOMIDATE 0.3 mg/kg IVP/IOP SINGLE DOSE

OR

PEDIATRIC: MIDAZOLAM 0.1 mg/kg TO MAX OF 5 mg IVP/IOP

FOR POST-ORAL INTUBATION SEDATION TO PREVENT EXTUBATION (IF INDICATED):

PEDIATRIC: MIDAZOLAM 0.1 mg/kg TO MAX OF 5 mg IVP/IOP

OR

PEDIATRIC: DIAZEPAM 0.1 mg/kg TO MAX OF 5 mg IVP/IOP

OR

PEDIATRIC: LORAZEPAM 0.1 mg/kg TO MAX OF 2 mg IVP/IOP

CONTINUOUS ASSESSMENT & TREATMENT PER APPLICABLE PROTOCOL(S)