



EMS System for Metropolitan Oklahoma City and Tulsa 2026 Medical Control Board Treatment Protocols



Approved 9/17/25, Effective 1/15/26, replaces all prior versions

Treatment Priorities

1. Oxygenation/Ventilation support

2G – MEDICATION ASSISTED INTUBATION

ADULT

PARAMEDIC

PARAMEDIC

MEDICATION ASSISTED INTUBATION IF INDICATED

FOLLOW PROTOCOL 2F – ORAL INTUBATION TECHNIQUE & CONFIRMATION OF INTUBATION

FOR FACILITATING ORAL INTUBATION:

ADULT: ETOMIDATE 0.3 mg/kg IVP/IOP SINGLE DOSE

OR

ADULT: MIDAZOLAM 0.1 mg/kg TO MAX OF 5mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP \geq 100 mmHg

OR

ADULT: KETAMINE 1.5 mg/kg TO MAX OF 200 mg IVP/IOP

FOR POST-ORAL INTUBATION SEDATION TO PREVENT EXTUBATION (IF INDICATED)

ADULT: MIDAZOLAM 0.1 mg/kg TO MAX OF 5mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP \geq 100 mmHg

OR

ADULT: DIAZEPAM 0.1 mg/kg TO MAX OF 5 mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP \geq 100 mmHg

OR

ADULT: LORAZEPAM 0.1 mg/kg TO MAX OF 2mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP \geq 100 mmHg

CONTINUOUS ASSESSMENT & TREATMENT PER APPLICABLE PROTOCOL(S)