Treatment Priorities

1. Oxygenation/Ventilation support

2G – MEDICATION ASSISTED INTUBATION

ADULT

PARAMEDIC

PARAMEDIC

MEDICATION ASSISTED INTUBATION IF INDICATED
FOLLOW PROTOCOL 2F – ORAL INTUBATION TECHNIQUE & CONFIRMATION OF INTUBATION

FOR FACILITATING ORAL INTUBATION:
ADULT: ETOMIDATE 0.3 mg/kg IVP/IOP SINGLE DOSE
OR
ADULT: MIDAZOLAM 0.1 mg/kg TO MAX OF 5mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP ≥100 mmHg
OR
ADULT: KETAMINE 1.5 mg/kg TO MAX OF 200 mg IVP/IOP

FOR POST-ORAL INTUBATION SEDATION TO PREVENT EXTBUATION (IF INDICATED)
ADULT: MIDAZOLAM 0.1 mg/kg TO MAX OF 5mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP ≥100 mmHg

OR
ADULT: DIAZEPAM 0.1 mg/kg TO MAX OF 5 mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP ≥100 mmHg

OR
ADULT: LORAZEPAM 0.1 mg/kg TO MAX OF 2mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP ≥100 mmHg

CONTINUOUS ASSESSMENT & TREATMENT PER APPLICABLE PROTOCOL(S)