



EMS System for Metropolitan Oklahoma City and Tulsa 2020 Medical Control Board Treatment Protocols



Approved 9/11/19, Effective 1/15/20, replaces all prior versions

2G - MEDICATION ASSISTED INTUBATION ADULT

TREATMENT PRIORITIES

1, Oxygenation/Ventilation support

EMERGENCY MEDICAL
DISPATCHER

EMERGENCY MEDICAL
RESPONDER

EMT

EMT-INTERMEDIATE 85

ADVANCED EMT

PARAMEDIC

PARAMEDIC

MEDICATION-ASSISTED INTUBATION IF INDICATED
FOLLOW PROTOCOL 2F – ORAL INTUBATION FOR TECHNIQUE & CONFIRMATION OF INTUBATION

FOR FACILITATING ORAL INTUBATION:

ADULT: ETOMIDATE 0.3 mg/kg IVP/IOP SINGLE DOSE
OR

ADULT: MIDAZOLAM 0.1 mg/kg TO MAX OF 5 mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP \geq 100 mmHg

FOR POST-ORAL INTUBATION SEDATION TO PREVENT EXTUBATION (IF INDICATED):

ADULT: MIDAZOLAM 0.1 mg/kg TO MAX OF 5 mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP \geq 100 mmHg
OR

ADULT: DIAZEPAM 0.1 mg/kg TO MAX OF 5 mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP \geq 100 mmHg
OR

ADULT: LORAZEPAM 0.1 mg/kg TO MAX OF 2 mg IVP/IOP
MAY REPEAT ONCE IF ADULT SYS BP \geq 100 mmHg

CONTINUOUS ASSESSMENT & TREATMENT PER APPLICABLE PROTOCOL(S)