



EMS System for Metropolitan Oklahoma City and Tulsa 2019 Medical Control Board Treatment Protocols



Approved 9/12/18, Effective 1/15/19

3M – DYSPNEA – CROUP PEDIATRIC

- TREATMENT PRIORITIES**
- Vital signs (including EtCO₂, if equipped)
 - Oxygenation support
 - O₂ by NC, NRB
 - BVM, ETT if indicated
 - Ventilation support
 - BVM, ETT if indicated
 - Nebulization therapy
 - Epinephrine 1mg/mL 1:1,000 at 3mL
 Consider Foreign body as a cause of stridor

EMD

ADVISE TO AVOID PHYSICAL EXERTION OR ENVIRONMENTAL STRESS (TEMP EXTREMES).
ADVISE PT SELF-ADMINISTRATION OF MEDICATIONS (eg. ALBUTEROL INHALER)
AS PREVIOUSLY PRESCRIBED FOR DYSPNEA SYMPTOMS

- EMERGENCY MEDICAL DISPATCHER
- EMERGENCY MEDICAL RESPONDER
- EMT
- EMT-INTERMEDIATE 85
- ADVANCED EMT
- PARAMEDIC

EMR | **EMT**

CROUP MOSTLY OCCURS IN INFANTS AND YOUNG CHILDREN BETWEEN SIX MONTHS AND THREE YEARS OF AGE, AND IS LESS COMMONLY SEEN IN CHILDREN OLDER THAN SIX YEARS

GENERAL SUPPORTIVE CARE
OBTAIN VITAL SIGNS
O₂ VIA NC, NRB, OR BVM AS APPROPRIATE
APPLY CARDIAC MONITOR (if equipped)
ATTEMPT TO KEEP CHILD CALM WHILE PROPERLY SECURING THE CHILD FOR TRANSPORT

EMT OR HIGHER LICENSE:
MEASURE END-TIDAL CO₂ & MONITOR WAVEFORM CAPNOGRAPHY (if equipped, **Mandatory use if pt intubated)

EMT-I85 | **AEMT**

PEDIATRIC: INTUBATE IF INDICATED PER PROTOCOL 17E

IV ACCESS
PEDIATRIC: IV NS TKO IF SYS BP ≥ (70 + 2x age in years) mmHg
PEDIATRIC: IV NS 20 mL/kg BOLUS IF SYS BP < (70 + 2x age in years) mmHg IF NO SIGNS OF PULMONARY EDEMA

PARAMEDIC

PEDIATRIC: METHYLPREDNISOLONE 2 mg/kg NOT TO EXCEED 125 mg IVP. MAY GIVE IM IF NO VASCULAR ACCESS OBTAINED.

FOR SIGNIFICANT INSPIRATORY STRIDOR AT REST, DECREASED RESPONSIVENESS, POOR PERFUSION, APNEA OR CYANOSIS
PEDIATRIC: NEBULIZED EPINEPHRINE 1mg/mL (1:1000) at 3mg/3mL VIA NEBULIZER

PEDIATRIC: MEDICATION-ASSISTED INTUBATION IF INDICATED PER PROTOCOL 17F
CONTINUOUS ASSESSMENT & TREATMENT PER APPLICABLE PROTOCOL(S)