



EMS System for Metropolitan Oklahoma City and Tulsa 2020 Medical Control Board Treatment Protocols



Approved 9/11/19, Effective 1/15/20, replaces all prior versions

4L- INTRA-ARREST WAKEFULNESS ADULT

TREATMENT PRIORITIES
 1. Safety of self
 2. Safety of public safety professionals
 3. Safety of patient
 4. Continuity of resuscitation

EMERGENCY MEDICAL DISPATCHER
EMERGENCY MEDICAL RESPONDER
EMT
EMT-INTERMEDIATE 85
ADVANCED EMT
PARAMEDIC

EMR	EMT
<p>ASSIST IN PHYSICAL CONTROL OF PATIENT, INCLUDING APPLYING PHYSICAL RESTRAINTS ANY RESTRAINT(S) SHOULD MINIMIZE ANY DETRIMENT TO RESPIRATORY OR PERFUSION MECHANICS</p> <p>USE ADEQUATE NUMBERS OF PUBLIC SAFETY PROFESSIONALS TO MINIMIZE RISK OF INJURY TO SELF AND OTHERS</p> <p>SPEAK CALMLY TO PATIENT WITH REASSURANCE THAT HELP IS BEING PROVIDED</p> <p>CONTINUE RESUSCITATION CARE PER APPLICABLE PROTOCOLS</p>	

EMT-I85	AEMT
<p>IV/IO ACCESS DO NOT RISK SELF INJURY WITH NEEDLESTICK IN IV ACCESS IF PT COMBATIVE</p>	

PARAMEDIC
<p>CHEMICAL RESTRAINT:</p> <p>ALL PATIENTS REQUIRING CHEMICAL RESTRAINT ARE TO BE PHYSICALLY RESTRAINED AS WELL</p> <p>ADULT: MIDAZOLAM 0.1 mg/kg IVP/IO TO MAX OF 5 mg. MAY REPEAT ONCE. OR ADULT: DIAZEPAM 5 mg IVP/IO IF MIDAZOLAM NOT AVAILABLE. MAY REPEAT ONCE. OR ADULT: LORAZEPAM 2 mg IVP/IO IF MIDAZOLAM NOT AVAILABLE. MAY REPEAT ONCE. (MIDAZOLAM STRONGLY PREFERRED DUE TO MOST RAPID ONSET OF ACTION OF BENZODIAZEPINE OPTIONS)</p> <p>CONTINUOUS ASSESSMENT & TREATMENT PER APPLICABLE PROTOCOL(S)</p>