



# EMS System for Metropolitan Oklahoma City and Tulsa 2026 Medical Control Board Treatment Protocols



Approved 9/17/25, Effective 1/15/26, replaces all prior versions

TREATMENT PRIORITIES
1. Vital signs
2. O <sub>2</sub>
3. Dextrose for hypoglycemia
4. Benzodiazepine for sustained, active seizure (refer to 6D Seizure if applicable)
5. BVM prior to administration of Naloxone
Evaluate differential diagnosis of Syncope & treat per protocol(s):
o Acute Coronary Syndrome
o Cardiac Dysrhythmia
o Hypotension (Shock)
o Hypoxemia (Shock)
o Head Injury
o Stroke
o Seizure
o Infection (Sepsis/ Meningitis)
o Medication/Alcohol
o Heat or Cold Illness
o Psychogenic/Emotion

6E - SYNCOPES ADULT & PEDIATRIC	
<b>EMD</b>	<b>EMD</b> KEEP PATIENT FREE FROM INJURY HAZARDS AVOID PLACING ANYTHING IN MOUTH ADVISE TO AVOID PHYSICAL EXERTION OR ENVIRONMENTAL STRESS (TEMP EXTREMES) PLACE IN RECOVERY POSITION/POSITION OF COMFORT
<b>EMR</b>	<b>EMT</b> GENERAL SUPPORTIVE CARE; OBTAIN VITAL SIGNS O <sub>2</sub> VIA NC, NRB, OR BVM AS APPROPRIATE DETERMINE BLOOD GLUCOSE FOR PATIENT ABLE TO SWALLOW <b>ADULT &amp; PEDIATRIC WEIGHT <math>\geq 25</math> kg HYPOGLYCEMIA CARE:</b> IF GLUCOSE <50 mg/dL, 1 tube ORAL GLUCOSE (15 grams) PO <b>PEDIATRIC WEIGHT &lt;25 kg HYPOGLYCEMIA CARE:</b> IF GLUCOSE <50 mg/dL, 1/2 tube ORAL GLUCOSE (7.5 grams) PO <b>TOXINS/DRUG OVERDOSE – SUSPECTED NARCOTIC/OPIATE</b> ADDRESS OXYGENATION AND VENTILATION (SP <sub>02</sub> $\geq 94\%$ ) BEFORE ADMINISTRATION OF NALOXONE <b>APNEIC/AGONALLY BREATHING</b> <b>ADULT:</b> NALOXONE 2 mg IN, MAY REPEAT ONCE <b>PEDIATRIC:</b> NALOXONE 0.5 mg IN, MAY REPEAT TO MAX OF 2 mg <b>INEFFECTIVE BREATHING ACTIVITY</b> <b>ADULT &amp; PEDIATRIC:</b> NALOXONE 0.5 mg IN, MAY REPEAT TO MAX OF 2 mg USE NALOXONE TO RESTORE EFFECTIVE BREATHING; AVOID EXCESSIVE DOSING TO PREVENT WITHDRAWAL  APPLY CARDIAC MONITOR/OBTAIN 12-LEAD ECG (if equipped) TRANSMIT 12-LEAD ECG TO RECEIVING EMERGENCY DEPARTMENT <b>EMT OR HIGHER LICENSE:</b> MEASURE END-TIDAL CO <sub>2</sub> & MONITOR WAVEFORM CAPNOGRAPHY (if equipped, *Mandatory use if pt intubated) PLACE SUPRAGLOTTIC AIRWAY IF INDICATED & ONLY IF BVM VENTILATIONS INEFFECTIVE
<b>EMT-I85</b> <b>AEMT</b>	

<b>EMERGENCY MEDICAL DISPATCHER</b>
<b>EMERGENCY MEDICAL RESPONDER</b>
<b>EMT</b>
<b>EMT-INTERMEDIATE 85</b>
<b>ADVANCED EMT</b>
<b>PARAMEDIC</b>

<b>EMT-I85</b> <b>AEMT</b>
IV ACCESS
<b>ADULT:</b> IV NS TKO IF SYS BP $\geq 100$ mmHg WITHOUT HYPOTENSIVE SYMPTOMS
<b>ADULT:</b> IV NS 250 mL BOLUS IF SYS BP <100 mmHg WITH HYPOTENSIVE SYMPTOMS & NO SIGNS OF PULMONARY EDEMA,
<b>ADULT:</b> REPEAT UP TO 2 LITERS NS IF SYS BP REMAINS < 100 mmHg WITH HYPOTENSIVE SYMPTOMS & NO SIGNS OF PULMONARY EDEMA
<b>PEDIATRIC:</b> IV NS TKO IF SYS BP $\geq (70 + 2 \times \text{age in years})$ mmHg
<b>PEDIATRIC:</b> IV NS 20 mL/kg BOLUS IF SYS BP < $(70 + 2 \times \text{age in years})$ mmHg IF NO SIGNS OF PULMONARY EDEMA
<b>HYPONATREMIA (GLUCOSE &lt;50 mg/dL) - ADULT &amp; PEDIATRIC:</b>
D10 5 mL/kg IVPB WIDE OPEN UP TO 250 mL OR
D25 2 mL/kg IV/IO UP TO 100 mL (must be $\geq 1$ year of age) OR D50 1 mL/kg IV/IO UP TO 50 mL (must be $\geq 25$ kg)
IF NO VASCULAR ACCESS OBTAINED & IF IO SEEMS EXCESSIVE TO CLINICAL STATUS:
GLUCAGON: IF PT WT $\geq 25$ kg, 1mg IM; <25 kg, 0.5 mg IM
<b>ADULT &amp; PEDIATRIC:</b> REPEAT DETERMINATION OF BLOOD GLUCOSE POST-HYPONATREMIA TREATMENT
<b>ADULT:</b> INTUBATE IF INDICATED; DO NOT INTUBATE PATIENTS WITH RAPIDLY REVERSIBLE AMS ETIOLOGY (eg. HYPOGLYCEMIA, OPIATES)
<b>ADVANCED EMT OR HIGHER LICENSE:</b>
<b>TOXINS/DRUG OVERDOSE – SUSPECTED NARCOTIC/OPIATE – APNEIC/AGONALLY BREATHING</b>
<b>ADULT:</b> NALOXONE 2 mg IVP/IO/IN MAY REPEAT ONCE
<b>PEDIATRIC:</b> NALOXONE 0.5 mg IVP/IO/IN, MAY REPEAT TO MAX OF 2 mg
<b>TOXINS/DRUG OVERDOSE – SUSPECTED NARCOTIC/OPIATE – INEFFECTIVE BREATHING ACTIVITY</b>
<b>ADULT &amp; PEDIATRIC:</b> NALOXONE 0.5 mg IVP/IO/IN, MAY REPEAT TO MAX OF 2 mg
USE NALOXONE TO RESTORE EFFECTIVE BREATHING; AVOID EXCESSIVE DOSING TO PREVENT WITHDRAWAL

<b>PARAMEDIC</b>
<b>ADULT:</b> MEDICATION-ASSISTED INTUBATION IF INDICATED CONTINUOUS ASSESSMENT & TREATMENT OF SUSPECTED AMS ETIOLOGY PER APPLICABLE PROTOCOL(S) CONSULT OLMC IF ABOVE TREATMENT INEFFECTIVE FOR HYPOGLYCEMIA OR NARCOTIC/OPIATE ETIOLOGY CONSULT OLMC IF UNCERTAIN OF ETIOLOGY AND TREATMENT PLAN OF AMS