



# EMS System for Metropolitan Oklahoma City and Tulsa 2019 Medical Control Board Treatment Protocols



Approved 3/13/19, Effective 6/1/19, replaces all prior versions

**TREATMENT PRIORITIES**

- Vital signs
- O<sub>2</sub>
- Dextrose for hypoglycemia
- Benzodiazepine for sustained, active seizure (refer to 6D Seizure if applicable)

Evaluate differential diagnosis of AMS & treat per protocol(s):

- Hypoxemia (Shock)
- Head Injury
- Stroke
- Seizure
- Infection (Sepsis/Meningitis)
- Medication/Alcohol
- Heat or Cold Illness

## 6B - ALTERED MENTAL STATUS ADULT & PEDIATRIC

**EMD**

KEEP PATIENT FREE FROM INJURY HAZARDS  
AVOID PLACING ANYTHING IN MOUTH  
PLACE IN RECOVERY POSITION POST SEIZURE

**EMERGENCY MEDICAL DISPATCHER**

**EMERGENCY MEDICAL RESPONDER**

**EMT**

**EMT-INTERMEDIATE 85**

**ADVANCED EMT**

**PARAMEDIC**

<b>EMR</b>	<b>EMT</b>
GENERAL SUPPORTIVE CARE & OBTAIN VITAL SIGNS O <sub>2</sub> VIA NC, NRB, OR BVM AS APPROPRIATE	
<b>TOXINS/DRUG OVERDOSE – SUSPECTED NARCOTIC/OPIATE APNEIC/AGONALLY BREATHING</b> <b>ADULT:</b> NALOXONE 2 mg IN, MAY REPEAT ONCE <b>PEDIATRIC:</b> NALOXONE 0.5 mg IN, MAY REPEAT TO MAX OF 2 mg	
<b>INEFFECTIVE BREATHING ACTIVITY</b> <b>ADULT &amp; PEDIATRIC:</b> NALOXONE 0.5 mg IN, MAY REPEAT TO MAX OF 2 mg USE NALOXONE TO RESTORE EFFECTIVE BREATHING; AVOID EXCESSIVE DOSING TO PREVENT WITHDRAWAL	
DETERMINE BLOOD GLUCOSE FOR PATIENT ABLE TO SWALLOW <b>ADULT &amp; PEDIATRIC WEIGHT ≥ 25 kg HYPOGLYCEMIA CARE:</b> IF GLUCOSE <50 mg/dL, 1 tube ORAL GLUCOSE (15 grams) PO <b>PEDIATRIC WEIGHT &lt; 25 kg HYPOGLYCEMIA CARE:</b> IF GLUCOSE <50 mg/dL, ½ tube ORAL GLUCOSE (7.5 grams) PO	
APPLY CARDIAC MONITOR (if equipped) <b>EMT OR HIGHER LICENSE:</b> MEASURE END-TIDAL CO <sub>2</sub> & MONITOR WAVEFORM CAPNOGRAPHY (if equipped, **Mandatory use if pt intubated) PLACE SUPRAGLOTTIC AIRWAY IF INDICATED & ONLY IF BVM VENTILATIONS INEFFECTIVE	

<b>EMT-I85</b>	<b>AEMT</b>
----------------	-------------

IV ACCESS  
**ADULT:** IV NS TKO IF SYS BP ≥ 100 mmHg WITHOUT HYPOTENSIVE SYMPTOMS  
**ADULT:** IV NS 250 mL BOLUS IF SYS BP <100 mmHg WITH HYPOTENSIVE SYMPTOMS & NO SIGNS OF PULMONARY EDEMA, REPEAT UP TO 2 LITERS NS IF SYS BP REMAINS < 100 mmHg WITH HYPOTENSIVE SYMPTOMS & NO SIGNS OF PULMONARY EDEMA  
**PEDIATRIC:** IV NS TKO IF SYS BP ≥ (70 + 2x age in years) mmHg  
**PEDIATRIC:** IV NS 20 mL/kg BOLUS IF SYS BP < (70 + 2x age in years) mmHg IF NO SIGNS OF PULMONARY EDEMA

**HYPOGLYCEMIA (GLUCOSE <50 mg/dL) - ADULT & PEDIATRIC**  
D10 5 mL/kg IVPB WIDE OPEN UP TO 250 mL OR  
D25 2 mL/kg IV/IO UP TO 100 mL (must be ≥ 1 year of age) OR D50 1 mL/kg IV/IO UP TO 50 mL (must be ≥ 25 kg)  
IF NO VASCULAR ACCESS OBTAINED & IF IO SEEMS EXCESSIVE TO CLINICAL STATUS:  
GLUCAGON: IF PT WT ≥25 kg, 1mg IM; <25 kg, 0.5 mg IM  
**ADULT & PEDIATRIC:** REPEAT DETERMINATION OF BLOOD GLUCOSE POST-HYPOGLYCEMIA TREATMENT

**ADULT:** INTUBATE IF INDICATED; DO NOT INTUBATE PATIENTS WITH RAPIDLY REVERSIBLE ETIOLOGY (eg. HYPOGLYCEMIA, OPIATES)

**ADVANCED EMT OR HIGHER LICENSE:**

**TOXINS/DRUG OVERDOSE – SUSPECTED NARCOTIC/OPIATE – APNEIC/AGONALLY BREATHING**  
**ADULT:** NALOXONE 2 mg IVP/IO/IN, MAY REPEAT ONCE  
**PEDIATRIC:** NALOXONE 0.5 mg IVP/IO/IN, MAY REPEAT TO MAX OF 2 mg

**TOXINS/DRUG OVERDOSE – SUSPECTED NARCOTIC/OPIATE – INEFFECTIVE BREATHING ACTIVITY**  
**ADULT & PEDIATRIC:** NALOXONE 0.5 mg IVP/IO/IN, MAY REPEAT TO MAX OF 2 mg  
USE NALOXONE TO RESTORE EFFECTIVE BREATHING; AVOID EXCESSIVE DOSING TO PREVENT WITHDRAWAL

**PARAMEDIC**

**ADULT:** MEDICATION-ASSISTED INTUBATION IF INDICATED  
CONTINUOUS ASSESSMENT & TREATMENT OF SUSPECTED AMS ETIOLOGY PER APPLICABLE PROTOCOL(S)  
CONSULT OLMC IF ABOVE TREATMENT INEFFECTIVE FOR HYPOGLYCEMIA OR NARCOTIC/OPIATE ETIOLOGY  
CONSULT OLMC IF UNCERTAIN OF ETIOLOGY AND TREATMENT PLAN OF AMS