


## EMS System for Metropolitan Oklahoma City and Tulsa 2025 Medical Control Board Treatment Protocols





Approved 9/04/24, Effective 1/15/25, replaces all prior versions



PEDIATRIC: ONDANSETRON 0.1 mg/kg IVP TO A MAXIMUM SINGLE DOSE OF 4 mg
IF AGE >2 years, MAY GIVE ONDANSETRON 4 mg ODT

SEPTIC SHOCK UNRESPONSIVE TO IV/IO FLUIDS AS ABOVE? **ADULT: PHARMACOLOGIC TREATMENT IF SYS BP < 100 mmHg:** NOREPINEPHRINE 2-4 mcg/min TITRATE TO SYS  $\geq$  100 mmHg **OR** 

DOPAMINE 5-20 mcg/kg/min TITRATE TO SYS BP ≥ 100 mmHg IF NOREPINEPHRINE NOT AVAILABLE PEDIATRIC: OLMC CONSULT FOR PHARMACOLOGIC TREATMENT

## FEBRILE SEIZURE?

IF SOLITARY AND SHORT DURATION (< 5 MINUTES) – OBSERVE IF PROLONGED/STATUS EPILEPTICUS, TREAT PER PROTOCOL 6D – SEIZURE

MINIMIZE SCENE TIME IF SEPSIS SUSPECTED

IF CLINICAL CONCERN FOR SEPSIS, NOTIFY RECEIVING ED OF POSSIBLE SEPSIS IN RADIO REPORT
CONTINUOUS ASSESSMENT & TREATMENT PER APPLICABLE PROTOCOL(S)