


## EMS System for Metropolitan Oklahoma City and Tulsa 2019 Medical Control Board Treatment Protocols





Approved 9/12/18, Effective 1/15/19, replaces all prior versions



ANTIEMETIC (IF ACTIVELY VOMITING) ADULT: ONDANSETRON 4 mg IVP/ODT. MAY REPEAT ONCE IN 10 MINUTES

PEDIATRIC: ONDANSETRON 0.1 mg/kg IVP TO A MAXIMUM SINGLE DOSE OF 4 mg
IF AGE >2 years, MAY GIVE ONDANSETRON 4 mg ODT

DOPAMINE 5-20 mcg/kg/min TITRATE TO SYS BP \(^2\) 100 mmHg IF NOREPINEPHRINE NOT AVAILABLE PEDIATRIC: OLMC CONSULT FOR PHARMACOLOGIC TREATMENT

## FEBRILE SEIZURE?

IF SOLITARY AND SHORT DURATION (< 5 MINUTES) – OBSERVE IF PROLONGED/STATUS EPILEPTICUS, TREAT PER PROTOCOL 6D – SEIZURE

MINIMIZE SCENE TIME IF SEPSIS SUSPECTED

IF CLINICAL CONCERN FOR SEPSIS, NOTIFY RECEIVING ED OF POSSIBLE SEPSIS IN RADIO REPORT CONTINUOUS ASSESSMENT & TREATMENT PER APPLICABLE PROTOCOL(S)